



Year: 2020

Nut consumption and the prevalence and severity of non-alcoholic fatty liver disease

Semmler, Georg ; Bachmayer, Sebastian ; Wernly, Sarah ; Wernly, Bernhard ; Niederseer, David ;
Huber-Schönauer, Ursula ; Stickel, Felix ; Aigner, Elmar ; Datz, Christian

Abstract: BACKGROUND Nut consumption has been associated with reduced inflammation, insulin resistance, and oxidative stress. However, the influence on the prevalence and severity of non-alcoholic fatty liver disease (NAFLD) has yet to be evaluated. METHODS 4655 subjects were included as part of a colorectal carcinoma screening program (SAKKOPI) between 07/2010 and 07/2019 and analyzed 2020. Patients were characterized using biochemical and metabolic parameters, as well as a detailed questionnaire on dietary habits. The diagnosis of NAFLD was established using abdominal ultrasound. Consumption of nuts was graded as: no consumption or <1 time/week, 1-6 times/week, 1 time/day and ≥ 2 times/day. RESULTS Mean age was 58.5 ± 9.8 years with a mean BMI of 26.5 ± 4.7 kg/m². 2058 (44.2%) patients suffered from the metabolic syndrome, 2407 (51.6%) had arterial hypertension, 2287 (49.1%) showed prediabetes/diabetes, 1854 (39.4%) had dyslipidemia and 1984 patients (43.5%) were diagnosed with NAFLD. Prevalence of metabolic syndrome (1219 [48.7%] vs. 605 [40.2%] vs. 189 [37.4%] vs. 45 [31.7%], $p < 0.001$) and NAFLD (1184 [48.1%] vs. 594 [40.7%] vs. 158 [31.7%] vs. 48 [34.0%], $p < 0.001$). On multivariable logistic regression analysis adjusting for potential confounders and dietary patterns, nut consumption ≥ 1 time/day was inversely associated with NAFLD in the overall cohort (adjusted Odds ratio[aOR]: 0.719 [95%CI: 0.558-0.926], $p = 0.011$). However, following subgroup analysis, this inverse association was only confirmed in male patients (aOR: 0.589 [95%CI: 0.411-0.844], $p = 0.004$) but not in females (aOR: 0.886 [95%CI: 0.616-1.275], $p = 0.515$). Moreover, patients who consumed nuts 1-6 times/week had a significantly lower prevalence of advanced fibrosis (Fib-4 score > 2.67 : aOR: 0.551 [95%CI: 0.338-0.898], $p = 0.017$; Forns-Index > 6.9 : aOR: 0.585 [95%CI: 0.402-0.850], $p = 0.005$). CONCLUSIONS Nut consumption might exert beneficial effects on the prevalence of NAFLD in males. The negative association with advanced fibrosis warrants further investigation.

DOI: <https://doi.org/10.1371/journal.pone.0244514>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-198582>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

Semmler, Georg; Bachmayer, Sebastian; Wernly, Sarah; Wernly, Bernhard; Niederseer, David; Huber-Schönauer, Ursula; Stickel, Felix; Aigner, Elmar; Datz, Christian (2020). Nut consumption and the prevalence and severity of non-alcoholic fatty liver disease. PLoS ONE, 15(12):e0244514.

DOI: <https://doi.org/10.1371/journal.pone.0244514>


RESEARCH ARTICLE

Nut consumption and the prevalence and severity of non-alcoholic fatty liver disease

Georg Semmler¹, Sebastian Bachmayer¹, Sarah Wernly¹, Bernhard Wernly², David Niederseer³, Ursula Huber-Schönauer¹, Felix Stickel⁴, Elmar Aigner⁵, Christian Datz^{1*}

1 Department of Internal Medicine, General Hospital Oberndorf, Teaching Hospital of the Paracelsus Medical University Salzburg, Oberndorf, Salzburg, Austria, **2** Second Department of Medicine, Paracelsus Medical University Salzburg, Salzburg, Austria, **3** Department of Cardiology, University Hospital Zurich, Zurich, Switzerland, **4** Department of Gastroenterology and Hepatology, University Hospital Zurich, Zurich, Switzerland, **5** First Department of Medicine, Paracelsus Medical University Salzburg, Salzburg, Austria

* c.datz@kh-oberndorf.at



Abstract

Background

Nut consumption has been associated with reduced inflammation, insulin resistance, and oxidative stress. However, the influence on the prevalence and severity of non-alcoholic fatty liver disease (NAFLD) has yet to be evaluated.

Methods

4655 subjects were included as part of a colorectal carcinoma screening program (SAK-KOPI) between 07/2010 and 07/2019 and analyzed 2020. Patients were characterized using biochemical and metabolic parameters, as well as a detailed questionnaire on dietary habits. The diagnosis of NAFLD was established using abdominal ultrasound. Consumption of nuts was graded as: no consumption or <1 time/week, 1–6 times/week, 1 time/day and ≥ 2 times/day.

Results

Mean age was 58.5 ± 9.8 years with a mean BMI of $26.5 \pm 4.7 \text{ kg/m}^2$. 2058 (44.2%) patients suffered from the metabolic syndrome, 2407 (51.6%) had arterial hypertension, 2287 (49.1%) showed prediabetes/diabetes, 1854 (39.4%) had dyslipidemia and 1984 patients (43.5%) were diagnosed with NAFLD. Prevalence of metabolic syndrome (1219 [48.7%] vs. 605 [40.2%] vs. 189 [37.4%] vs. 45 [31.7%], $p < 0.001$) and NAFLD (1184 [48.1%] vs. 594 [40.7%] vs. 158 [31.7%] vs. 48 [34.0%], $p < 0.001$). On multivariable logistic regression analysis adjusting for potential confounders and dietary patterns, nut consumption ≥ 1 time/day was inversely associated with NAFLD in the overall cohort (adjusted Odds ratio[aOR]: 0.719 [95%CI: 0.558–0.926], $p = 0.011$). However, following subgroup analysis, this inverse association was only confirmed in male patients (aOR: 0.589 [95%CI: 0.411–0.844], $p = 0.004$) but not in females (aOR: 0.886 [95%CI: 0.616–1.275], $p = 0.515$). Moreover, patients who consumed nuts 1–6 times/week had a significantly lower prevalence of advanced fibrosis

OPEN ACCESS

Citation: Semmler G, Bachmayer S, Wernly S, Wernly B, Niederseer D, Huber-Schönauer U, et al. (2020) Nut consumption and the prevalence and severity of non-alcoholic fatty liver disease. PLoS ONE 15(12): e0244514. <https://doi.org/10.1371/journal.pone.0244514>

Editor: Pavel Strnad, Medizinische Fakultät der RWTH Aachen, GERMANY

Received: October 15, 2020

Accepted: December 10, 2020

Published: December 31, 2020

Copyright: © 2020 Semmler et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Data cannot be shared publicly because of institutional data policy restrictions and national data policy restrictions (www.ris.bka.gv.at). Data are available from the General Hospital Oberndorf (contact via [w.ferlitz@kh-oberndorf.at](mailto:ferlitz@kh-oberndorf.at)) for researchers who meet the criteria for access to confidential data.

Funding: No funding was received for this study. SPAR Österreichische Warenhandels-AG did not provide financial support in the form of authors' salaries and/or research materials, and did not play

any role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: All authors declare that they have no conflicts of interest regarding the submitted work. The following authors disclose conflicts of interests outside the submitted work: C. D. is a member of the scientific advisory board of SPAR Österreichische Warenhandels-AG. This does not alter our adherence to PLOS ONE policies on sharing data and materials.

Abbreviations: BMI, body mass index; BP, blood pressure; CCS, chronic coronary syndrome; FFQ, food frequency questionnaire; HDL-C, high-density lipoprotein cholesterol; HOMA-IR, Homeostasis Model Assessment for Insulin Resistance; IFG, impaired fasting glucose; IGT, impaired glucose tolerance; IQR, interquartile range; IR, insulin resistance; LDL-C, low-density lipoprotein cholesterol; MetS, metabolic syndrome; NAFLD, non-alcoholic fatty liver disease; NFS, NAFLD fibrosis score; OGTT, oral glucose tolerance test; PAD, peripheral artery disease; SSB, sugar-sweetened beverage; T2DM, type 2 diabetes mellitus; WC, waist circumference.

(Fib-4 score >2.67: aOR: 0.551 [95%CI: 0.338–0.898], $p = 0.017$; Forns-Index >6.9: aOR: 0.585 [95%CI: 0.402–0.850], $p = 0.005$).

Conclusions

Nut consumption might exert beneficial effects on the prevalence of NAFLD in males. The negative association with advanced fibrosis warrants further investigation.

Introduction

Non-alcoholic fatty liver disease (NAFLD) is regarded the emerging complex chronic liver disease of the 21st century, being significantly associated with increased cardiovascular and liver-related risk [1–3]. Despite the increasing prevalence of NAFLD, no medical therapy has been established. Thus, lifestyle and dietary interventions remain the cornerstone of treatment in these patients.

Nuts are nutrient-dense food containing bioactive compounds regarded beneficial for health: vegetable protein, unsaturated fatty acids, fiber, minerals, vitamins, tocopherols, phytosterols, and polyphenols [4]. However, each nut type has a unique composition and different amounts led to different health outcomes in interventional studies [5]. Importantly, most data supporting a beneficial effect of nuts exist for almonds, walnuts, pistachios, and peanuts, followed other nut types [5]. For instance, two randomized controlled trials could confirm the lipid-lowering effect of 10/60g almonds/day in patients with coronary artery disease or type 2 diabetes mellitus (T2DM) [6, 7]. In large population-based studies not differentiating between nut types [4], their consumption has been associated with reduced inflammation [8, 9] and insulin resistance (IR) [10]. Moreover, a large body of evidence exists for the protective effect of nuts on T2DM [11, 12], metabolic syndrome (MetS) [12–14], obesity [10, 12, 13] and arterial hypertension [15]. Apart from these beneficial effects on aspects of the MetS, their role in chronic liver disease and especially NAFLD merits further investigation since inflammation, oxidative stress and IR are considered key drivers of NAFLD [16–18]. Recently, a Chinese study reported a significantly lower prevalence of NAFLD in patients consuming nuts ≥ 4 times/week [19]. However, another Chinese study confirmed this inverse association of NAFLD and nut consumption only in a small subgroup of men when consuming ≥ 8.86 g/day [20] while data from Caucasian subjects addressing this topic is scarce. Therefore, we aimed to further elucidate the role of nut consumption on the prevalence and severity of NAFLD.

Materials & methods

Patients and design

4655 individuals participating in a screening program for colorectal cancer (SAKKOPI) between 07/2010 and 07/2019 were included in this analysis. Only patients with established liver disease (i.e. viral hepatitis, autoimmune hepatitis, hereditary liver diseases) or with consumption of a significant amount of alcohol (≥ 2 drinks/day for females and ≥ 3 drinks/day for males) were excluded from the analysis. All other patients without established liver disease were included portraying a representative sample of the general population. Patients were carefully characterized including laboratory and clinical data, and completed a detailed questionnaire on lifestyle and dietary habits. The study design and details of the clinical and biochemical work-up of included subjects have been reported previously [21]. The study was

approved by the “Ethikkommission für das Bundesland Salzburg” (approval no. 415-E/1262/2-2010), and all patients gave written informed consent to participate.

Dietary assessment

Dietary data were collected using a food-frequency questionnaire (FFQ) which consisted of 69-items and was conducted in line with the questionnaire used in the SAPHIR [22], as well as the EPIC study [23–25]. Patients completed the FFQ self-administered prior to all examinations. Data on alcohol consumption, intake of fast-food, vegetables, fruits, sweets, red and processed meat, white meat, fish, coffee and consumption of sugar-sweetened beverages (SSB) and nuts were assessed. Specifically, nut consumption was reported semiquantitatively as no consumption or <1 time (corresponding to ~10g) per week, 1–6 times per week, 1 time per day and ≥ 2 times per day. Different types of nuts were not specified, and legume seeds were included according to the interpretation of the patient. In order to quantify healthy eating patterns, an adopted version of the “Alternate Healthy Eating Index” (AHEI-2010) was calculated in accordance with the FFQ being available [26].

Definitions

The diagnosis of NAFLD was established using ultrasound being performed by experienced operators. The liver was considered “normal” if the echogenicity was homogenous and similar or slightly higher than that of the renal parenchyma. Portal veins and diaphragm had to be visualized. The liver was considered as “fatty liver” when areas of significant increased echogenicity in relation to the renal parenchyma were found. The severity of sonographic steatosis was not graded [21]. Advanced fibrosis ($\geq F3$) was assessed using Fib-4 score, NAFLD fibrosis score (NFS) and Forns-Index [27, 28]. Cut-offs for advanced fibrosis were determined as previously described: >2.67 [27, 29] for Fib-4 score, >0.676 for NFS [30] and >6.9 for Forns-Index [31]. Metabolic characterization included an oral glucose tolerance test (OGTT) as well as measurement of fasting blood glucose and insulin measurement to assess IR. T2DM was defined as either blood glucose level of ≥ 200 mg/dl after 2 hours following oral glucose tolerance test (OGTT), fasting blood glucose (FBG) ≥ 125 mg/dl, HbA1c $\geq 6.5\%$, or previously prescribed antidiabetic medication including insulin. Impaired fasting glucose (IFG) was defined as FBG 100–124mg/dl in non-diabetic individuals. Impaired glucose tolerance (IGT) was defined as a blood glucose of 140–199mg/dl after 2 hours following OGTT in non-diabetic individuals. Prediabetes was defined as IFG or IGT in non-diabetic individuals. Dysglycemia was defined as presence of either prediabetes or T2DM. Components of the MetS were defined according to the revised definition of the International Diabetes Federation (2005) [32]. Visceral obesity was defined as waist circumference (WC) ≥ 80 cm for women and ≥ 94 cm for men. Hypertension was defined as a blood pressure (BP) $\geq 130/85$ mmHg or previous prescription of any antihypertensive drug. Dyslipidemia was defined as high-density lipoprotein cholesterol (HDL-C) <40 mg/dl in males or <50 mg/dl in females, triglycerides ≥ 150 mg/dl or previously prescribed lipid lowering drugs. The MetS was defined as at least three of the following components: Visceral adiposity, hypertension, FBG ≥ 100 mg/dl or antidiabetic therapy, HDL-C <40 mg/dl in males or <50 mg/dl in females or antilipidemic drugs, triglycerides ≥ 150 mg/dl or antilipidemic drugs. Of note, levels of systolic BP, FBG, blood glucose following OGTT, HbA1c, HOMA-IR, triglycerides, cholesterol, HDL-C and low-density lipoprotein cholesterol (LDL-C) were only considered for comparison in the absence of specific medication. Chronic coronary syndrome (CCS) was defined as history of myocardial infarction, coronary artery disease, coronary artery bypass graft or coronary stent.

Statistical analyses

Statistical analyses were performed using IBM SPSS Statistics 26 (SPSS Inc., Armonk, New York, USA) and GraphPad Prism 8 (GraphPad Software, La Jolla, California, USA) in 2020. Continuous variables were reported as mean \pm standard deviation or median (interquartile range [IQR]) depending on their distribution, while categorical variables were shown as numbers and proportions of patients (%). Comparisons of continuous variables were performed using Student's t-test, Mann-Whitney U test or Kruskal-Wallis test, as applicable. Group comparisons were performed applying Chi-squared test. Logistic regression analysis was performed to investigate the associations of nut consumption on the presence of NAFLD and the presence of advanced fibrosis, using the group with the lowest nut consumption as reference category. Since the number of individuals in group 4 was too limited to draw firm conclusions, groups 3 and 4 were combined for logistic regression analyses. P-values were calculated for each group compared to the reference group, as well as P-values for linear trends. Multivariable models were adjusted for potential confounders including sex, age and BMI, MetS, and dietary parameters such as alcohol consumption, intake of fast-food, vegetables, fruits, sweets, red and processed meat, white meat, fish, coffee and consumption of SSB, as previously described [19, 20]. Analyses on Fib-4 score and Forns-Index were not adjusted for age since this variable is included in these indices. Subgroup analyses were performed in patients with NAFLD, male and female individuals. A two-sided p-value ≤ 0.05 was considered as statistically significant.

Results

Patient characteristics

Overall, 4655 individuals were included with a slight preponderance of males ($n = 2395$, 51.5%), and a mean age of 58.5 ± 9.8 . 2058 patients (44.2%) suffered from the MetS with 3278 (76.9%) suffering from visceral obesity, 2407 (51.6%) from hypertension, 2287 (49.1%) from T2DM or prediabetes, and 1854 (39.8%) from dyslipidemia. Additionally, 280 (6.0%) individuals had previously diagnosed CCS, 152 (3.3%) peripheral arterial disease (PAD) and 148 (3.2%) had previously suffered from stroke. 2502 patients (53.7%) did not consume nuts or consumed nuts < 1 time/week, 1506 patients (32.4%) consumed nuts 1–6 times/week, 505 patients (10.8%) consumed nuts 1 time/day (once daily) and 142 patients (3.1%) ≥ 2 times/day (at least twice daily).

Nut consumption, metabolic syndrome and cardiovascular health

The prevalence of the MetS declined with more frequent nut consumption (48.7% vs. 40.2% vs. 37.4% vs. 31.7%, $p < 0.001$). Importantly, this trend was also observed for all singular components (Table 1). Similarly, the proportion of patients suffering from cardiovascular diseases decreased with increasing nut consumption: 7.5% vs. 4.4% vs. 4.0% vs. 3.5% for CCS ($p < 0.001$), 4.3% vs. 2.0% vs. 2.6% vs. 2.1% for PAD ($p = 0.001$) and 4.0% vs. 2.1% vs. 2.8% vs. 1.4% for stroke ($p = 0.005$).

Dietary patterns including consumption of alcohol, fast-food, fruits, vegetables, sweets, coffee, SSBs, red meat and fish were significantly different among these groups with less frequent fast-food consumption and consumption of red or processed meat and a higher consumption of fruits, vegetables and coffee in patients with more frequent nut consumption (Table 2). Consecutively, these subjects had higher scores in the adopted version of the AHEI-2010 (42.4 ± 8.7 vs. 49.3 ± 8.2 vs. 57.6 ± 7.8 vs. 60.6 ± 8.2 points, $p < 0.001$).

Table 1. Patient characteristics in the overall cohort and compared among patients grouped according to their frequency of nut consumption.

Patient characteristics	All patients, n = 4655	<1 time/week, n = 2502	1–6 times/week, n = 1506	1 time/day, n = 505	≥2 times/day, n = 142	P value
Age, years	58.5±9.8	58.8±10.2	57.6±9.2	59.8±8.7	58.2±8.8	<0.001
Male sex	2395 (51.5%)	1360 (54.4%)	766 (50.9%)	209 (41.4%)	60 (42.3%)	<0.001
Obesity	1068 (22.9%)	661 (26.5%)	302 (20.1%)	77 (15.2%)	29 (19.7%)	<0.001
BMI, kg/mg ²	26.5±4.7	27.6±4.9	26.7±4.5	25.9±4.2	25.9±4.5	<0.001
Visceral obesity*	3278 (76.9%)	1814 (79.9%)	1050 (75.1%)	331 (71.3%)	83 (62.4%)	<0.001
WC, cm*	96.3±13.3	98.1±13.6	95.2±12.6	92.5±12.3	92.2±14.2	<0.001
Metabolic syndrome	2058 (44.2%)	1219 (48.7%)	605 (40.2%)	189 (37.4%)	45 (31.7%)	<0.001
Hypertension	2407 (51.6%)	1776 (71.0%)	972 (64.5%)	341 (67.5%)	94 (66.2%)	<0.001
T2DM/prediabetes	2287 (49.1%)	1334 (53.3%)	679 (45.1%)	226 (44.8%)	48 (33.8%)	<0.001
T2DM	638 (13.7%)	422 (16.9%)	151 (10.0%)	47 (9.3%)	18 (12.7%)	<0.001
Prediabetes	1649 (35.4%)	912 (36.5%)	528 (35.1%)	179 (35.4%)	30 (21.1%)	0.003
IFG	1365 (29.3%)	753 (30.1%)	446 (29.6%)	139 (27.5%)	27 (19.0%)	0.001
IGT	638 (13.7%)	362 (14.5%)	193 (12.8%)	76 (15.0%)	7 (4.9%)	<0.001
FBG, mg/dl	98±20	99±15	98±27	96±11	96±18	0.008
OGTT after 2h, mg/dl	122±37	125±39	119±33	122±33	119±39	<0.001
HbA1c, %	5.5±0.4	5.6±0.5	5.5 ±0.4	5.5±0.4	5.5±0.4	<0.001
HOMA-IR, points	1.67 (1.11–2.62)	1.79 (1.15–2.81)	1.63 (1.13–2.53)	1.46 (1.00–2.30)	1.46 (0.90–2.30)	<0.001
Dyslipidemia	1854 (39.8%)	1083 (43.3%)	553 (36.7%)	174 (14.5%)	44 (31.0%)	<0.001
Triglycerides, mg/dl	102 (76–142)	106 (78–148)	101 (74–140)	93 (72–129)	88 (67–126)	<0.001
Hypertriglyceridemia	879 (18.9%)	507 (20.3%)	281 (18.7%)	72 (14.3%)	19 (13.4%)	<0.001
Cholesterol, mg/dl	224±41	223±43	226±40	225±39	222±41	0.333
Hypercholesterinemia	2857 (61.4%)	1463 (58.5%)	991 (65.8%)	315 (62.4%)	88 (62.0%)	0.040
HDL-C, mg/dl	60±17	59±17	60±18	62±17	64±17	<0.001
LDL-C, mg/dl	145±37	145±38	147±37	145±36	140±37	0.244
CCS	280 (6.0%)	188 (7.5%)	67 (4.4%)	20 (4.0%)	5 (3.5%)	<0.001
PAD	152 (3.3%)	106 (4.3%)	30 (2.0%)	13 (2.6%)	3 (2.1%)	0.001
Stroke	148 (3.2%)	100 (4.0%)	32 (2.1%)	14 (2.8%)	2 (1.4%)	0.005

* Data available in 4265 patients.

Abbreviations: BMI–body mass index; CCS–chronic coronary syndrome; FBG–fasting blood glucose; HOMA-IR–Homeostasis Model Assessment for Insulin Resistance; IFG–impaired fasting glucose; IGT–impaired glucose tolerance; OGTT–oral glucose tolerance test; PAD–peripheral artery disease; T2DM–type 2 diabetes mellitus; WC–waist circumference.

<https://doi.org/10.1371/journal.pone.0244514.t001>

Nut consumption and NAFLD

The prevalence of NAFLD decreased among nut consumption groups: 1184 patients (48.1%) with nut consumption <1time/week, 594 patients (40.7%) with 1–6 times/week, 158 (31.7%) with 1time/day and 48 patients (34.0%) with ≥2times/day were diagnosed with NAFLD ($p<0.001$, Table 3). The prevalence of a Fib-4 score of >2.67 indicative for advanced fibrosis was significantly different among groups (3.9% vs. 2.0% vs. 3.2% vs. 3.5%, $p = 0.014$), concordantly to the prevalence of NFS >0.676 (5.6% vs. 1.9% vs. 3.0% vs. 2.4%, $p<0.001$) and Forns-Index >6.9 (6.9% vs. 3.4% vs. 4.2% vs. 2.1%, $p<0.001$). Finally, patients had less elevated levels of gamma-glutamyl transferase (GGT; median GGT/sex-specific upper limit of normal: 0.69 [IQR: 0.46–1.18] vs. 0.64 [0.44–1.08] vs. 0.56 [0.38–0.87] vs. 0.51 [0.37–0.82], $p<0.001$).

On multivariable logistic regression analysis adjusting for potential confounders, patients consuming nuts ≥1time/day had a significantly lower risk for NAFLD (aOR: 0.589 [95%CI: 0.558–0.926], $p = 0.011$) when compared to those within the lowest group (Table 4). However, a linear trend across all groups was observed ($p = 0.016$). Interestingly, patients consuming

Table 2. Dietary patterns compared among patients grouped according to their frequency of nut consumption.

Patient characteristics	<1 time/week, n = 2502	1–6 times/week, n = 1506	1 time/day, n = 505	≥2 times/day, n = 142	P value
Alcohol consumption ≥3 times/week	31.0%	37.5%	29.7%	32.4%	<0.001
Fast food ≥1/month	51.0%	57.9%	41.5%	34.8%	<0.001
Daily fruit intake	59.5%	71.8%	84.5%	84.1%	<0.001
Daily vegetable intake	63.8%	75.4%	86.2%	86.3%	<0.001
Daily sweets intake	30.8%	37.3%	41.5%	39.1%	<0.001
Daily coffee intake	48.6%	58.4%	53.1%	54.9%	<0.001
Daily SSB intake	10.2%	13.9%	10.3%	9.9%	0.002
Red meat ≥2days/week	71.3%	65.4%	55.7%	46.6%	<0.001
White meat ≥2days/week	32.5%	32.8%	31.9%	27.5%	0.663
Fish ≥1day/week	76.4%	86.8%	83.8%	85.5%	<0.001
Adopted AHEI-2010*	42.4±8.7	49.3±8.2	57.6±7.8	60.6±8.2	<0.001

* The percentage of trans-Isomers of fatty acids and polyunsaturated fatty acids from total energy intake could not be sufficiently calculated owing to the nature of our FFQ. Therefore, these variables could not be calculated leading to a maximum score of 90 points.

Abbreviations: SSB=sugar-sweetened beverages.

<https://doi.org/10.1371/journal.pone.0244514.t002>

nuts 1–6 times/week had a significantly lower odds of advanced fibrosis (Fib-4 score >2.67: aOR: 0.551 [95%CI: 0.338–0.898], $p = 0.017$; Forns-Index >6.9: aOR: 0.585 [95%CI: 0.402–0.850], $p = 0.005$) which was more pronounced in the subgroup of NAFLD patients (Fib-4 score >2.67: aOR: 0.275 [95%CI: 0.112–0.678], $p = 0.005$; Forns-Index >6.9: aOR: 0.384 [95%CI: 0.226–0.653], $p < 0.001$). In order to identify potential reasons for these inverse associations, markers of inflammation were compared among subgroups of nut-consumption. Interestingly, subjects with more frequent nut consumption had significantly lower levels of C-reactive protein concentrations (0.20 [IQR: 0.10–0.40] vs. 0.17 [0.10–0.31] vs. 0.15 [0.08–0.30] vs. 0.11 [0.08–0.27] mg/dl, $p < 0.001$) and ferritin concentrations (134 [71–231] vs. 114 [63–193] vs. 103 [54–178] vs. 89 [42–159] µg/l, $p < 0.001$; S1 Table in [S1 File](#)). Of note, when

Table 3. Distribution of hepatic steatosis as well as non-invasive scores for hepatic fibrosis compared among patients grouped according to their frequency of nut consumption.

Patient characteristics	<1 time/week, n = 2502	1–6 times/week, n = 1506	1 time/day, n = 505	≥2 times/day, n = 142	P value
NAFLD*	1184 (48.1%)	594 (40.7%)	158 (31.7%)	48 (34.0%)	<0.001
Fib-4 score	1.16 (0.89–1.53)	1.12 (0.88–1.43)	1.17 (0.95–1.58)	1.19 (1.00–1.54)	<0.001
>2.67	96 (3.9%)	30 (2.0%)	16 (3.2%)	5 (3.5%)	0.014
>3.25	58 (2.3%)	20 (1.3%)	10 (2.0%)	1 (0.7%)	0.108
NFS	-1.64 (-2.41–[-0.70])	-1.88 (-2.51–[-1.17])	-1.81 (-2.49–[-0.93])	-1.81 (-2.30–[-0.83])	<0.001
>0.676	64 (5.6%)	17 (1.9%)	9 (3.0%)	2 (2.4%)	<0.001
Forns-Index	4.51±1.55	4.29±1.36	4.37±1.37	4.31±1.31	<0.001
>6.9	171 (6.9%)	51 (3.4%)	21 (4.2%)	3 (2.1%)	<0.001
ALT/ULN	0.50 (0.38–0.71)	0.50 (0.37–0.69)	0.49 (0.37–0.66)	0.49 (0.37–0.58)	0.055
AST/ULN	0.60 (0.51–0.74)	0.60 (0.49–0.71)	0.57 (0.49–0.71)	0.60 (0.51–0.66)	0.056
GGT/ULN	0.69 (0.46–1.18)	0.64 (0.44–1.08)	0.56 (0.38–0.87)	0.51 (0.37–0.82)	<0.001

* Data available in 4563 patients.

ULN was regarded as 50 U/L for ALT and AST in male patients and 35 U/L in female patients.

Abbreviations: ALT=alanine aminotransferase; AST=aspartate aminotransferase; GGT=gamma-glutamyl transferase; NAFLD=non-alcoholic fatty liver disease; NFS=NAFLD fibrosis score; ULN=upper limit of normal.

<https://doi.org/10.1371/journal.pone.0244514.t003>

Table 4. Odds Ratio (OR) and 95% confidence interval (95%CI) for NAFLD and advance fibrosis among groups of nut-consumers using binary logistic regression analyses adjusted for potential confounders.

		Overall cohort			NAFLD		
		aOR (95% CI)	P value	P value for linear trend	aOR (95% CI)	P value	P value for linear trend
NAFLD	<1 time/week	reference	reference	0.016	-	-	-
	1–6 times/week	0.923 (0.770–1.106)	0.384		-	-	-
	≥1 time/day	0.719 (0.558–0.926)	0.011		-	-	-
Fib-4 score >2.67	<1 time/week	reference	reference	0.110	reference	reference	0.099
	1–6 times/week	0.551 (0.338–0.898)	0.017		0.275 (0.112–0.678)	0.005	
	≥1 time/day	0.780 (0.427–1.423)	0.418		0.844 (0.337–2.110)	0.716	
Forns-Index >6.9	<1 time/week	reference	reference	0.002	reference	reference	0.001
	1–6 times/week	0.585 (0.402–0.850)	0.005		0.384 (0.226–0.653)	<0.001	
	≥1 time/day	0.533 (0.304–0.934)	0.028		0.450 (0.198–1.023)	0.057	

Analyses were based on the overall cohort and NAFLD patients.

Displayed OR are adjusted for sex, age, BMI, metabolic syndrome, hepatic steatosis, alcohol consumption, intake of fast-food, vegetables, fruits, sweets, red and processed meat, white meat, fish, coffee and consumption of SSB. OR for Fib-4 score and Forns-Index were not adjusted for age since this variable is included in these indices.

Abbreviations: aOR—adjusted Odds ratio; NAFLD—non-alcoholic fatty liver disease; SSB—sugar-sweetened beverage; T2DM—type 2 diabetes mellitus; 95%CI— 95% confidence interval.

<https://doi.org/10.1371/journal.pone.0244514.t004>

replacing dietary patterns with the adopted AHEI-2010 for regression analyses, there was no independently association with NAFLD (aOR per point: 0.992 [95%CI: 0.984–1.000], $p = 0.063$).

Gender differences

In order to address gender differences with regard to nut consumption, we performed gender-specific subgroup analyses in males and females. Differences in baseline characteristics are displayed in S2 Table in [S1 File](#). The negative association with NAFLD was even more pronounced in males consuming nuts ≥ 1 time/day (aOR: 0.589 [95%CI: 0.411–0.844], $p = 0.004$). However, it could not be confirmed in female patients (aOR: 0.886 [95%CI: 0.616–1.275], $p = 0.516$; S3 Table in [S1 File](#)). Although a trend for decreased odds of having advanced fibrosis as assessed by Fib-4 score >2.67 was evident both in males and females consuming nuts 1–6 times/week, a significant associated could only be confirmed in males applying the Forns-index >6.9 (aOR: 0.610 [95%CI: 0.402–0.924], $p = 0.020$). Again, the adopted AHEI-2010 was not associated with NAFLD in males (aOR: 0.992, [95%CI: 0.982–1.003], $p = 0.156$) or females (aOR: 0.994, [95%CI: 0.982–1.007], $p = 0.347$).

Discussion

Nuts have previously been associated with a reduced risk for MetS, especially T2DM, hypertension and obesity [11–14], and reduced risk for cardiovascular disease and death [33], while recently, a potentially protective effect on the prevalence of NAFLD was proposed in two Asian cohorts [19, 20]. In our study, we show that nut consumption at least once a day was associated with a significantly decreased risk for NAFLD while any consumption of nuts was linked to a lower prevalence of advanced fibrosis in NAFLD subjects. Moreover, we confirm other study results showing a lower prevalence of MetS and cardiovascular diseases among nut-consumers [13, 14].

Generally speaking, nuts are rich in monounsaturated and polyunsaturated fatty acids (PUFA), vegetable protein, fibers, folate, vitamins (e.g. tocopherols) and minerals (e.g. copper),

among other bioactive compounds including carotenoids, phytosterols and polyphenolic compounds including flavonoids [5, 34–36]. Specifically, folate has been associated with counter-acting atherothrombotic properties of homocysteine [37] while phytosterols might interfere with cholesterol absorption [4]. Tocopherol (i.e. vitamin E) and polyphenols have protective properties against free radicals and oxidative stress [4, 38], with flavonoid intake being linked to reduced all-cause, cardiovascular and cancer mortality [39], and tocopherol being widely used off-label to treat NAFLD due to its beneficial effects on liver enzyme elevations and histology improvement [40]. Other important components are omega-3-fatty-acids: Their dietary supplementation has recently been associated with reduced triglyceride levels and cardioprotective effects [41, 42]. Carotenoids—especially β -carotene—have also been attributed certain antioxidative effects, and recent study results showed that consuming high amounts of carotenoids lowered the risk of NAFLD [43], and improved lipid profiles in an animal models [44]. Finally, nondigestible components of nuts act as probiotics providing beneficial substrates for the gut microbiota [45]. Thus, several components and their synergetic interaction could explain a potential favorable effect on human physiology [4]. Interestingly, copper deficiency has previously been linked to disturbed iron homeostasis in NAFLD [46], potentially aggravating the natural course of NAFLD [47, 48]. Although copper supplementation by nut consumption might counteract harmful effects of copper deficiency in NAFLD, considerable variations in copper content of different nuts need to be considered.

Recently, Zhang et al [19] reported on a large retrospective cohort of 23915 patients excluding those with cardiovascular diseases or cancer, and found a significantly lower prevalence of NAFLD in patients consuming nuts ≥ 4 times/week, being equivalent to ≥ 200 –240g/week. However, a smaller Chinese cohort study including 1068 patients, could only reproduce this finding in males within the highest quartile of nut intake [20]. Together, these studies add up to evidence from small cohort studies (< 400 patients) reporting benefits for subjects with NAFLD [49–51]. Although we confirm these data in a representative cohort of Caucasian individuals, subgroup analyses could only reproduce this inverse association in male individuals. This is interesting since our findings go in line with previous studies reporting an association only in males [20, 49]. However, we can only speculate on potential explanations: First, the benefit of nut consumption might be better discernible in men on the background of more prevalent NAFLD and more pronounced MetS components [52]. Moreover, men usually have larger visceral fat mass, being associated with a more pro-inflammatory profile [53], which directly releases free-fatty acids into the portal vein, thereby promoting fat-accumulation in the liver [54]. Second, women are considered to have a higher awareness for healthy nutrition [55], which is also indicated by a higher prevalence of female gender in more frequent nut consumption groups in our study. Thus, the effect of nuts could be mitigated by other healthy dietary patterns.

To the best of our knowledge, this is the first study which proposes a beneficial effect of nuts on the prevalence of advanced fibrosis in patients with NAFLD. Although we cannot explain why the effect on fibrosis seems to start already when consuming nuts less frequently (i.e. 1–6 times/week), our data allow speculation that the beneficial effect on subsequent inflammation in the liver might already be evident at lower levels. Of note, we show that inflammation markers gradually decline with more frequent nut consumption, presumably indicating less inflammation, being the most important driver of fibrogenesis [56]. However, we acknowledge that blood inflammation markers may not fully mirror inflammation in the liver.

Additionally, it is unclear whether the beneficial effect of nuts follows a linear or U-shaped curve. Despite the large number of favorable bioactive compounds, nuts are also rich in fat and calories [4], consequently boosting one's energy intake if consumed on a daily basis. This

is supported by similar cholesterol and LDL-C levels among our patient groups independent from their amount of nut consumption. Thus, a frequent consumption might in some terms limit the beneficial effect of nuts' healthy components.

This study has strengths with regard to investigating the link between fatty liver and dietary habits. To begin with, patients underwent careful metabolic and characterization in the absence of clinical suspicion or indication for diagnostic work-up. Due to the coincidence that endoscopies are mainly performed at the General Hospital Oberndorf in the local catchment area, we are confident that we could obtain a representative sample of the general population at risk for NAFLD (~60 years of age). Importantly, the fact that no other regulations existed on which patients were included in the study (e.g. type of insurance or comorbidities), mitigated selection bias, which is often introduced in other liver-centered studies.

However, several limitations need to be considered when interpreting our results. To begin with, we acknowledge that nut types are unique in their composition and their effect on human health [5] especially when prepared differently (e.g. roasted, cooked, salted). However, we did not specifically assess nut types and preparation. Thus, we cannot exclude that differences in preparation (e.g. with chocolate in candy or baked goods) mitigate the associations reported in this study, and might decrease the study's impact. This is especially true for peanuts, which are commonly misclassified as nuts despite being legumes, which is likely to have occurred in our study. However, from the patient's point-of-view, specifying different nut types would increase uncertainties on true consumption since their frequency again decreases, and overreporting. From a statistical point of view, it would decrease statistical power. Therefore, we could not dissect potentially diverse effects of different nut types. Secondly, dietary habits assessed using FFQ are always patient-reported and therefore subject to response and recall bias: The correctness of the answers given by the patients can never be verified, nor can the correct recall from the patients' memory be verified. Also, FFQ require literacy and can be subject to overreporting [57].

Data from our study indicate that nuts are part of a healthier dietary pattern with significant differences among several healthy or unhealthy food types (Table 2). Demonstrated by a higher score in the AHEI-2020, these patients adhere to a healthier diet, which could accentuate the associations which were attributed to nut consumption. Nevertheless, the inverse association of nut consumption with NAFLD and advanced fibrosis remained significant after adjusting for several confounding food types such as red meat, coffee, and SSB, which have been linked to NAFLD previously [58–62]. Noteworthy, these significant dietary differences between patients consuming nuts and those who do not can also be observed in the majority of studies investigating the role of nuts [14], especially in those investigating NAFLD [19, 20].

Specific limitations of this study include the retrospective design which can only generate associations, but never causalities. Unfortunately, physical activity was only inconsistently reported and therefore could not be included as a covariate in regression models. Finally, NAFLD was diagnosed by abdominal ultrasound in our study. This is a limitation since hepatic steatosis can only be graded qualitatively, with particular weakness for lower degrees of liver fat. Therefore, a modest effect of any nut consumption might not be detected as a resolution of hepatic steatosis in our study.

Nevertheless, we demonstrate an inverse association of nut consumption with the prevalence of NAFLD in a large European cohort. In line with the literature, we confirm a stronger association in male gender. Moreover, we are the first to report a potentially favorable effect on the prevalence of advanced fibrosis. Our findings support recommendations for regular nut consumption as part of a Mediterranean diet, especially in patients with NAFLD [63]. Future dietary interventional studies investigating lipid-lowering and anti-inflammatory

properties of nuts in patients with NAFLD are needed to investigate the full potential of this unique food type as an add-on treatment of NAFLD.

Supporting information

S1 File.
(DOCX)

Author Contributions

Conceptualization: Georg Semmler, Christian Datz.

Data curation: Georg Semmler, Sebastian Bachmayer, Sarah Wernly, Ursula Huber-Schönauer, Christian Datz.

Formal analysis: Georg Semmler, David Niederseer, Felix Stickel, Elmar Aigner, Christian Datz.

Investigation: Georg Semmler, Christian Datz.

Methodology: Georg Semmler, Bernhard Wernly, Christian Datz.

Project administration: Christian Datz.

Resources: Christian Datz.

Writing – original draft: Georg Semmler, Christian Datz.

Writing – review & editing: Georg Semmler, Sebastian Bachmayer, Sarah Wernly, Bernhard Wernly, David Niederseer, Ursula Huber-Schönauer, Felix Stickel, Elmar Aigner, Christian Datz.

References

1. Younossi ZM, Marchesini G, Pinto-Cortez H, Petta S. Epidemiology of Nonalcoholic Fatty Liver Disease and Nonalcoholic Steatohepatitis: Implications for Liver Transplantation. *Transplantation*. 2019; 103(1):22–7. <https://doi.org/10.1097/TP.0000000000002484> PMID: 30335697
2. Niederseer D, Wernly S, Bachmayer S, Wernly B, Bakula A, Huber-Schönauer U, et al. Diagnosis of Non-Alcoholic Fatty Liver Disease (NAFLD) Is Independently Associated with Cardiovascular Risk in a Large Austrian Screening Cohort. *J Clin Med*. 2020; 9(4):1065.
3. Henson JB, Simon TG, Kaplan A, Osganian S, Masia R, Corey KE. Advanced fibrosis is associated with incident cardiovascular disease in patients with non-alcoholic fatty liver disease. *Alimentary pharmacology & therapeutics*. 2020; 51(7):728–36. <https://doi.org/10.1111/apt.15660> PMID: 32043602
4. Ros E. Health Benefits of Nut Consumption. *Nutrients*. 2010; 2(7):652–82. <https://doi.org/10.3390/nu2070652> PMID: 22254047
5. de Souza RGM, Schincaglia RM, Pimentel GD, Mota JF. Nuts and Human Health Outcomes: A Systematic Review. *Nutrients*. 2017; 9(12). <https://doi.org/10.3390/nu9121311> PMID: 29207471
6. Li SC, Liu YH, Liu JF, Chang WH, Chen CM, Chen CY. Almond consumption improved glycemic control and lipid profiles in patients with type 2 diabetes mellitus. *Metabolism: clinical and experimental*. 2011; 60(4):474–9. <https://doi.org/10.1016/j.metabol.2010.04.009> PMID: 20580779
7. Jamshed H, Sultan FA, Iqbal R, Gilani AH. Dietary Almonds Increase Serum HDL Cholesterol in Coronary Artery Disease Patients in a Randomized Controlled Trial. *J Nutr*. 2015; 145(10):2287–92. <https://doi.org/10.3945/jn.114.207944> PMID: 26269239
8. Yu Z, Malik VS, Keum N, Hu FB, Giovannucci EL, Stampfer MJ, et al. Associations between nut consumption and inflammatory biomarkers. *Am J Clin Nutr*. 2016; 104(3):722–8. <https://doi.org/10.3945/ajcn.116.134205> PMID: 27465378
9. Jiang R, Jacobs DR Jr., Mayer-Davis E, Szklo M, Herrington D, Jenny NS, et al. Nut and seed consumption and inflammatory markers in the multi-ethnic study of atherosclerosis. *American journal of epidemiology*. 2006; 163(3):222–31. <https://doi.org/10.1093/aje/kwj033> PMID: 16357111

10. Casas-Agustench P, Bullo M, Ros E, Basora J, Salas-Salvado J. Cross-sectional association of nut intake with adiposity in a Mediterranean population. *Nutrition, metabolism, and cardiovascular diseases: NMCD*. 2011; 21(7):518–25. <https://doi.org/10.1016/j.numecd.2009.11.010> PMID: 20219336
11. Jiang R, Manson JE, Stampfer MJ, Liu S, Willett WC, Hu FB. Nut and peanut butter consumption and risk of type 2 diabetes in women. *Jama*. 2002; 288(20):2554–60. <https://doi.org/10.1001/jama.288.20.2554> PMID: 12444862
12. Ibarrola-Jurado N, Bullo M, Guasch-Ferre M, Ros E, Martinez-Gonzalez MA, Corella D, et al. Cross-sectional assessment of nut consumption and obesity, metabolic syndrome and other cardiometabolic risk factors: the PREDIMED study. *PLoS One*. 2013; 8(2):e57367. <https://doi.org/10.1371/journal.pone.0057367> PMID: 23460844
13. Li H, Li X, Yuan S, Jin Y, Lu J. Nut consumption and risk of metabolic syndrome and overweight/obesity: a meta-analysis of prospective cohort studies and randomized trials. *Nutrition & metabolism*. 2018; 15:46. <https://doi.org/10.1186/s12986-018-0282-y> PMID: 29977320
14. Zhang Y, Zhang DZ. Relationship Between Nut Consumption and Metabolic Syndrome: A Meta-Analysis of Observational Studies. *Journal of the American College of Nutrition*. 2019; 38(6):499–505. <https://doi.org/10.1080/07315724.2018.1561341> PMID: 30716015
15. Djousse L, Rudich T, Gaziano JM. Nut consumption and risk of hypertension in US male physicians. *Clinical nutrition (Edinburgh, Scotland)*. 2009; 28(1):10–4. <https://doi.org/10.1016/j.clnu.2008.08.005> PMID: 18834651
16. Gao B, Tsukamoto H. Inflammation in Alcoholic and Nonalcoholic Fatty Liver Disease: Friend or Foe? *Gastroenterology*. 2016; 150(8):1704–9. <https://doi.org/10.1053/j.gastro.2016.01.025> PMID: 26826669
17. Kitade H, Chen G, Ni Y, Ota T. Nonalcoholic Fatty Liver Disease and Insulin Resistance: New Insights and Potential New Treatments. *Nutrients*. 2017; 9(4). <https://doi.org/10.3390/nu9040387> PMID: 28420094
18. Polimeni L, Del Ben M, Baratta F, Perri L, Albanese F, Pastori D, et al. Oxidative stress: New insights on the association of non-alcoholic fatty liver disease and atherosclerosis. *World journal of hepatology*. 2015; 7(10):1325–36. <https://doi.org/10.4254/wjh.v7.i10.1325> PMID: 26052378
19. Zhang S, Fu J, Zhang Q, Liu L, Meng G, Yao Z, et al. Association between nut consumption and non-alcoholic fatty liver disease in adults. *Liver international: official journal of the International Association for the Study of the Liver*. 2019; 39(9):1732–41. <https://doi.org/10.1111/liv.14164> PMID: 31162803
20. Bb Chen, Han Y, Pan X, Yan J, Liu W, Li Y, et al. Association between nut intake and non-alcoholic fatty liver disease risk: a retrospective case-control study in a sample of Chinese Han adults. *BMJ Open*. 2019; 9(9):e028961. <https://doi.org/10.1136/bmjopen-2019-028961> PMID: 31488477
21. Stadlmayr A, Aigner E, Steger B, Scharinger L, Lederer D, Mayr A, et al. Nonalcoholic fatty liver disease: an independent risk factor for colorectal neoplasia. *Journal of internal medicine*. 2011; 270(1):41–9. <https://doi.org/10.1111/j.1365-2796.2011.02377.x> PMID: 21414047
22. Kiechl S, Pechlaner R, Willeit P, Notdurfter M, Paulweber B, Willeit K, et al. Higher spermidine intake is linked to lower mortality: a prospective population-based study. *Am J Clin Nutr*. 2018; 108(2):371–80. <https://doi.org/10.1093/ajcn/nqy102> PMID: 29955838
23. Bohlscheid-Thomas S, Hoting I, Boeing H, Wahrendorf J. Reproducibility and relative validity of food group intake in a food frequency questionnaire developed for the German part of the EPIC project. *European Prospective Investigation into Cancer and Nutrition. International journal of epidemiology*. 1997; 26 Suppl 1:S59–70.
24. Bohlscheid-Thomas S, Hoting I, Boeing H, Wahrendorf J. Reproducibility and relative validity of energy and macronutrient intake of a food frequency questionnaire developed for the German part of the EPIC project. *European Prospective Investigation into Cancer and Nutrition. International journal of epidemiology*. 1997; 26 Suppl 1:S71–81. https://doi.org/10.1093/ije/26.suppl_1.s71 PMID: 9126535
25. Boeing H, Wahrendorf J, Becker N. EPIC-Germany—A source for studies into diet and risk of chronic diseases. *European Investigation into Cancer and Nutrition. Annals of nutrition & metabolism*. 1999; 43(4):195–204. <https://doi.org/10.1159/000012786> PMID: 10592368
26. Chiuve SE, Fung TT, Rimm EB, Hu FB, McCullough ML, Wang M, et al. Alternative Dietary Indices Both Strongly Predict Risk of Chronic Disease. *The Journal of Nutrition*. 2012; 142(6):1009–18. <https://doi.org/10.3945/jn.111.157222> PMID: 22513989
27. Tsochatzis EA, Newsome PN. Non-alcoholic fatty liver disease and the interface between primary and secondary care. *The Lancet Gastroenterology & Hepatology*. 2018; 3(7):509–17. [https://doi.org/10.1016/S2468-1253\(18\)30077-3](https://doi.org/10.1016/S2468-1253(18)30077-3) PMID: 29893235
28. Hagstrom H, Talback M, Andreasson A, Walldius G, Hammar N. Ability of Noninvasive Scoring Systems to Identify Individuals in the Population at Risk for Severe Liver Disease. *Gastroenterology*. 2020; 158(1):200–14. <https://doi.org/10.1053/j.gastro.2019.09.008> PMID: 31563624

29. McPherson S, Hardy T, Dufour JF, Petta S, Romero-Gomez M, Allison M, et al. Age as a Confounding Factor for the Accurate Non-Invasive Diagnosis of Advanced NAFLD Fibrosis. *The American journal of gastroenterology*. 2017; 112(5):740–51. <https://doi.org/10.1038/ajg.2016.453> PMID: [27725647](#)
30. Angulo P, Hui JM, Marchesini G, Bugianesi E, George J, Farrell GC, et al. The NAFLD fibrosis score: a noninvasive system that identifies liver fibrosis in patients with NAFLD. *Hepatology (Baltimore, Md)*. 2007; 45(4):846–54. <https://doi.org/10.1002/hep.21496> PMID: [17393509](#)
31. Forns X, Ampurdanès S, Llovet JM, Aponte J, Quintó L, Martínez-Bauer E, et al. Identification of chronic hepatitis C patients without hepatic fibrosis by a simple predictive model. *Hepatology (Baltimore, Md)*. 2002; 36(4 Pt 1):986–92. <https://doi.org/10.1053/jhep.2002.36128> PMID: [12297848](#)
32. Alberti KG, Eckel RH, Grundy SM, Zimmet PZ, Cleeman JI, Donato KA, et al. Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation*. 2009; 120(16):1640–5. <https://doi.org/10.1161/CIRCULATIONAHA.109.192644> PMID: [19805654](#)
33. Guasch-Ferré M, Liu X, Malik VS, Sun Q, Willett WC, Manson JE, et al. Nut Consumption and Risk of Cardiovascular Disease. *Journal of the American College of Cardiology*. 2017; 70(20):2519–32. <https://doi.org/10.1016/j.jacc.2017.09.035> PMID: [29145952](#)
34. Souza RG, Gomes AC, Naves MM, Mota JF. Nuts and legume seeds for cardiovascular risk reduction: scientific evidence and mechanisms of action. *Nutrition reviews*. 2015; 73(6):335–47. <https://doi.org/10.1093/nutrit/nuu008> PMID: [26011909](#)
35. Bolling BW, McKay DL, Blumberg JB. The phytochemical composition and antioxidant actions of tree nuts. *Asia Pacific journal of clinical nutrition*. 2010; 19(1):117–23. PMID: [20199996](#)
36. Kornsteiner M, Wagner K-H, Elmadfa I. Tocopherols and total phenolics in 10 different nut types. *Food Chemistry*. 2006; 98(2):381–7.
37. Welch GN, Loscalzo J. Homocysteine and atherothrombosis. *The New England journal of medicine*. 1998; 338(15):1042–50. <https://doi.org/10.1056/NEJM199804093381507> PMID: [9535670](#)
38. Taş NG, Gökmen V. Phenolic compounds in natural and roasted nuts and their skins: a brief review. *Current Opinion in Food Science*. 2017; 14:103–9.
39. Bondonno NP, Dalggaard F, Kyrø C, Murray K, Bondonno CP, Lewis JR, et al. Flavonoid intake is associated with lower mortality in the Danish Diet Cancer and Health Cohort. *Nature Communications*. 2019; 10(1):3651. <https://doi.org/10.1038/s41467-019-11622-x> PMID: [31409784](#)
40. Sanyal AJ, Chalasani N, Kowdley KV, McCullough A, Diehl AM, Bass NM, et al. Pioglitazone, Vitamin E, or Placebo for Nonalcoholic Steatohepatitis. *New England Journal of Medicine*. 2010; 362(18):1675–85.
41. Bhatt DL, Steg PG, Miller M, Brinton EA, Jacobson TA, Ketchum SB, et al. Cardiovascular Risk Reduction with Icosapent Ethyl for Hypertriglyceridemia. *New England Journal of Medicine*. 2018; 380(1):11–22. <https://doi.org/10.1056/NEJMoa1812792> PMID: [30415628](#)
42. Skulas-Ray AC, Wilson PWF, Harris WS, Brinton EA, Kris-Etherton PM, Richter CK, et al. Omega-3 Fatty Acids for the Management of Hypertriglyceridemia: A Science Advisory From the American Heart Association. *Circulation*. 2019; 140(12):e673–e91. <https://doi.org/10.1161/CIR.0000000000000709> PMID: [31422671](#)
43. Christensen K, Lawler T, Mares J. Dietary Carotenoids and Non-Alcoholic Fatty Liver Disease among US Adults, NHANES 2003(–)2014. *Nutrients*. 2019; 11(5).
44. Elvira-Torales LI, Martin-Pozuelo G, Gonzalez-Barrio R, Navarro-Gonzalez I, Pallares FJ, Santaella M, et al. Ameliorative Effect of Spinach on Non-Alcoholic Fatty Liver Disease Induced in Rats by a High-Fat Diet. *International journal of molecular sciences*. 2019; 20(7). <https://doi.org/10.3390/ijms20071662> PMID: [30987167](#)
45. Lamuel-Raventos RM, Onge MS. Prebiotic nut compounds and human microbiota. *Critical reviews in food science and nutrition*. 2017; 57(14):3154–63. <https://doi.org/10.1080/10408398.2015.1096763> PMID: [27224877](#)
46. Aigner E, Theurl I, Haufe H, Seifert M, Hohla F, Scharinger L, et al. Copper availability contributes to iron perturbations in human nonalcoholic fatty liver disease. *Gastroenterology*. 2008; 135(2):680–8. <https://doi.org/10.1053/j.gastro.2008.04.007> PMID: [18505688](#)
47. Datz C, Müller E, Aigner E. Iron overload and non-alcoholic fatty liver disease. *Minerva endocrinologica*. 2017; 42(2):173–83. <https://doi.org/10.23736/S0391-1977.16.02565-7> PMID: [27834478](#)
48. Aigner E, Strasser M, Haufe H, Sonnweber T, Hohla F, Stadlmayr A, et al. A role for low hepatic copper concentrations in nonalcoholic Fatty liver disease. *The American journal of gastroenterology*. 2010; 105(9):1978–85. <https://doi.org/10.1038/ajg.2010.170> PMID: [20407430](#)

49. Han JM, Jo AN, Lee SM, Bae HS, Jun DW, Cho YK, et al. Associations between intakes of individual nutrients or whole food groups and non-alcoholic fatty liver disease among Korean adults. *Journal of gastroenterology and hepatology*. 2014; 29(6):1265–72. <https://doi.org/10.1111/jgh.12520> PMID: [24955455](https://pubmed.ncbi.nlm.nih.gov/24955455/)
50. Kalafati IP, Dimitriou M, Borsa D, Vlachogiannakos J, Revenas K, Kokkinos A, et al. Fish intake interacts with TM6SF2 gene variant to affect NAFLD risk: results of a case-control study. *European journal of nutrition*. 2019; 58(4):1463–73. <https://doi.org/10.1007/s00394-018-1675-4> PMID: [29574608](https://pubmed.ncbi.nlm.nih.gov/29574608/)
51. Cueto-Galán R, Barón FJ, Valdivielso P, Pintó X, Corbella E, Gómez-Gracia E, et al. Changes in fatty liver index after consuming a Mediterranean diet: 6-year follow-up of the PREDIMED-Malaga trial. *Medicina clinica*. 2017; 148(10):435–43. <https://doi.org/10.1016/j.medcli.2016.11.032> PMID: [28126231](https://pubmed.ncbi.nlm.nih.gov/28126231/)
52. Lazo M, Hernaez R, Eberhardt MS, Bonekamp S, Kamel I, Guallar E, et al. Prevalence of nonalcoholic fatty liver disease in the United States: the Third National Health and Nutrition Examination Survey, 1988–1994. *American journal of epidemiology*. 2013; 178(1):38–45. <https://doi.org/10.1093/aje/kws448> PMID: [23703888](https://pubmed.ncbi.nlm.nih.gov/23703888/)
53. Ballestri S, Nascimbeni F, Baldelli E, Marrazzo A, Romagnoli D, Lonardo A. NAFLD as a Sexual Dimorphic Disease: Role of Gender and Reproductive Status in the Development and Progression of Non-alcoholic Fatty Liver Disease and Inherent Cardiovascular Risk. *Advances in therapy*. 2017; 34(6):1291–326. <https://doi.org/10.1007/s12325-017-0556-1> PMID: [28526997](https://pubmed.ncbi.nlm.nih.gov/28526997/)
54. Nielsen S, Guo Z, Johnson CM, Hensrud DD, Jensen MD. Splanchnic lipolysis in human obesity. *J Clin Invest*. 2004; 113(11):1582–8. <https://doi.org/10.1172/JCI21047> PMID: [15173884](https://pubmed.ncbi.nlm.nih.gov/15173884/)
55. Kiefer I, Rathmann T, Kunze M. Eating and dieting differences in men and women. *The Journal of Men's Health & Gender*. 2005; 2(2):194–201.
56. Koyama Y, Brenner DA. Liver inflammation and fibrosis. *J Clin Invest*. 2017; 127(1):55–64. <https://doi.org/10.1172/JCI88881> PMID: [28045404](https://pubmed.ncbi.nlm.nih.gov/28045404/)
57. Steinemann N, Grize L, Ziesemer K, Kauf P, Probst-Hensch N, Brombach C. Relative validation of a food frequency questionnaire to estimate food intake in an adult population. *Food & nutrition research*. 2017; 61(1):1305193.
58. Zelber-Sagi S, Ivancovsky-Wajcman D, Fliss Isakov N, Webb M, Orenstein D, Shibolet O, et al. High red and processed meat consumption is associated with non-alcoholic fatty liver disease and insulin resistance. *Journal of hepatology*. 2018; 68(6):1239–46. <https://doi.org/10.1016/j.jhep.2018.01.015> PMID: [29571924](https://pubmed.ncbi.nlm.nih.gov/29571924/)
59. Chen YP, Lu FB, Hu YB, Xu LM, Zheng MH, Hu ED. A systematic review and a dose-response meta-analysis of coffee dose and nonalcoholic fatty liver disease. *Clinical nutrition (Edinburgh, Scotland)*. 2019; 38(6):2552–7. <https://doi.org/10.1016/j.clnu.2018.11.030> PMID: [30573353](https://pubmed.ncbi.nlm.nih.gov/30573353/)
60. Asgari-Taei F, Zerafati-Shoae N, Dehghani M, Sadeghi M, Baradaran HR, Jazayeri S. Association of sugar sweetened beverages consumption with non-alcoholic fatty liver disease: a systematic review and meta-analysis. *European journal of nutrition*. 2019; 58(5):1759–69. <https://doi.org/10.1007/s00394-018-1711-4> PMID: [29761318](https://pubmed.ncbi.nlm.nih.gov/29761318/)
61. Alferink LJM, Fittipaldi J, Kieffe-de Jong JC, Taimr P, Hansen BE, Metselaar HJ, et al. Coffee and herbal tea consumption is associated with lower liver stiffness in the general population: The Rotterdam study. *Journal of hepatology*. 2017; 67(2):339–48. <https://doi.org/10.1016/j.jhep.2017.03.013> PMID: [28578837](https://pubmed.ncbi.nlm.nih.gov/28578837/)
62. Alferink LJM, Kieffe-de Jong JC, Erler NS, de Kneegt RJ, Hoorn EJ, Ikram MA, et al. Diet-dependent acid load—the missing link between an animal protein-rich diet and non-alcoholic fatty liver disease? *The Journal of clinical endocrinology and metabolism*. 2019.
63. Zelber-Sagi S, Salomone F, Mlynarsky L. The Mediterranean dietary pattern as the diet of choice for non-alcoholic fatty liver disease: Evidence and plausible mechanisms. *Liver International*. 2017; 37(7):936–49. <https://doi.org/10.1111/liv.13435> PMID: [28371239](https://pubmed.ncbi.nlm.nih.gov/28371239/)